



## **TheracosBio Extends Partnership with Mark Cuban Cost Plus Drug Company**

***BRENZAVVY Is Now Available to Healthcare Businesses via Cost Plus Drugs Marketplace at [business.costplusdrugs.com](https://business.costplusdrugs.com)***

**Marlborough, MA – February 14, 2024** – [TheracosBio](https://www.theracosbio.com) today announced that it is extending its collaboration with the Mark Cuban Cost Plus Drug Company. Under the partnership, BRENZAVVY® (bexagliflozin) is available directly to healthcare businesses under the Cost Plus Drugs Marketplace program.

BRENZAVVY represents a new approach to diabetes care. It is the first affordably priced, once-daily SGLT2 inhibitor, indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes.

The low cost of BRENZAVVY – the list price is 90% below other medicines in its class -- allows for expanded access for the uninsured and improves care for those with insurance by eliminating the hassles of insurance-company utilization management and cost-sharing. Pharmacies, hospitals and other healthcare businesses can now purchase the medicine through a Cost Plus Drugs site designed for businesses -- [business.costplusdrugs.com](https://business.costplusdrugs.com) -- for \$47.85 per each 30 tablet bottle.

“We launched BRENZAVVY in 2023 by making it available directly to patients at a transparent, low price through Mark Cuban Cost Plus Drugs online pharmacy and have been thrilled with the results,” said Brian Connelly, President and CEO of TheracosBio. “Through the new Cost Plus Drugs Marketplace, we are excited to provide affordably priced BRENZAVVY to healthcare businesses who are on the frontline in helping their patients and customers manage diabetes. This expansion of our collaboration should

make it easier than ever for patients to access BRENZAVVY, whether online or through their healthcare provider, clinic, or pharmacy.”

“Mark Cuban Cost Plus Drug Company has recently expanded its mission to make medications more affordable for healthcare businesses with the launch of the Cost Plus Drugs Marketplace,” said Dr. Alex Oshmyansky, CEO and Co-founder of Mark Cuban Cost Plus Drug Company. “This new platform provides healthcare businesses, including pharmacies, hospitals, surgery centers, and other healthcare providers, with access to a wide array of prescription and over-the-counter medications. Sourced from NABP-accredited suppliers, these drugs are offered at competitive prices to ensure that healthcare businesses can procure essential medications cost-effectively.”

Approximately 1 in 10 adults in the United States have type 2 diabetes, but suboptimal adherence to medication is common. A study of adherence among adults in the Medicare Part D database, published in the journal *Clinical Therapeutics*, estimated that 35.1% of patients were non-adherent to oral hypoglycemic agents. Cost-related nonadherence was reported by 17.6% of diabetic adults in the 18- to 64-year-old age group in a 2022 study published in *Diabetes Care*.

BRENZAVVY is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes. BRENZAVVY is not recommended for use to improve glycemic control in patients with type 1 diabetes mellitus. Phase 3 clinical studies have shown BRENZAVVY significantly reduces hemoglobin A1c and fasting blood sugar after 24 weeks, either as a monotherapy, in combination with metformin, or as an add-on to standard-of-care treatment consisting of a variety of regimens, including metformin, sulfonylureas, insulin, DPP4 inhibitors or combinations of these agents. Although BRENZAVVY is not approved for weight or blood pressure reduction, modest decreases in both weight and blood pressure have been observed in the clinical program.

BRENZAVVY is available as 20 mg oral tablets recommended to be taken once daily, in the morning with or without food. BRENZAVVY is

contraindicated in patients who are hypersensitive to bexagliflozin or any other ingredient in the BRENZAVVY tablet. The most common side effects of taking BRENZAVVY include female genital mycotic infections, urinary tract infections, and changes in urination, but more serious side effects are possible (see below, Important Safety Information about BRENZAVVY).

BRENZAVVY treatment can be initiated in adults with type 2 diabetes with an estimated glomerular filtration rate (eGFR) greater than or equal to 30 mL/min/1.73 m<sup>2</sup>. Patients with eGFR between 30 and 59 mL/min/1.73 m<sup>2</sup> are said to be in stage 3 chronic kidney disease. BRENZAVVY was the first SGLT2 inhibitor shown in a randomized, controlled clinical trial to be effective in adults with type 2 diabetes and stage 3 (3a + 3b) chronic kidney disease.

To learn more about BRENZAVVY and for full prescribing information visit <https://www.brenzavvy.com>. To report suspected adverse reactions, contact TheracosBio at 1-855-BRENZAVVY (1-855-273-6928) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### **Important Safety Information about BRENZAVVY**

**Limitation of Use:** BRENZAVVY (bexagliflozin) is not recommended for use to improve glycemic control in patients with type 1 diabetes mellitus. It may increase the risk of diabetic ketoacidosis in these patients.

### **Contraindications**

BRENZAVVY is contraindicated in patients with hypersensitivity to bexagliflozin or any excipient in BRENZAVVY. Anaphylaxis and angioedema have been reported with sodium-glucose co-transporter 2 (SGLT2) inhibitors.

### **Warnings and Precautions**

#### **Diabetic Ketoacidosis in Patients with Type 1 Diabetes Mellitus and Other Ketoacidosis**

BRENZAVVY increases the risk of life-threatening ketoacidosis in patients with type 1 diabetes. Type 2 diabetes and pancreatic disorders are also risk factors for ketoacidosis and fatal events of ketoacidosis have been reported

in patients with type 2 diabetes using SGLT2 inhibitors. Precipitating conditions for diabetic ketoacidosis or other ketoacidosis include under-insulinization due to insulin dose reduction or missed insulin doses, acute febrile illness, reduced caloric intake, ketogenic diet, surgery, volume depletion, and alcohol abuse. Signs and symptoms of diabetic ketoacidosis are consistent with dehydration and severe metabolic acidosis and include nausea, vomiting, abdominal pain, generalized malaise, and shortness of breath. Assess patients who present with signs and symptoms of metabolic ketoacidosis, regardless of blood glucose levels. If suspected, discontinue BRENZAVVY, treat promptly and monitor for resolution before restarting. Consider ketone monitoring in patients with type 1 diabetes mellitus as well as in others at risk for ketoacidosis. Withhold BRENZAVVY in clinical situations known to predispose to ketoacidosis and resume when clinically stable. Educate all patients on the signs and symptoms of ketoacidosis and instruct patients to discontinue BRENZAVVY and seek medical attention immediately if signs and symptoms occur.

### **Lower Limb Amputation**

Lower limb amputations have been observed in patients treated with BRENZAVVY in a study of patients with type 2 diabetes who had either established cardiovascular disease (CVD) or were at risk for CVD. Of the 23 BRENZAVVY-treated patients who had amputations, 15 were amputations of the toe and midfoot and 8 were amputations above and below the knee. Some patients had multiple amputations. Lower limb infections, gangrene, ischemia, and osteomyelitis were the most common precipitating medical events leading to the need for an amputation. The risk of amputation was highest in patients with a baseline history of prior amputation, peripheral vascular disease, and neuropathy.

Before initiating BRENZAVVY, consider factors in the patient's history that may predispose to the need for amputations, such as a history of prior amputation, peripheral vascular disease, neuropathy, and diabetic foot ulcers. Counsel patients receiving BRENZAVVY about the importance of routine preventative foot care and monitor for signs and symptoms of diabetic foot infection (including osteomyelitis), new pain or tenderness, sores or ulcers involving the lower limbs, and institute appropriate treatment.

### **Volume Depletion**

BRENZAVVY can cause intravascular volume contraction which may sometimes manifest as symptomatic hypotension or acute transient changes in creatinine. Acute kidney injury requiring hospitalization and dialysis has been reported in patients with type 2 diabetes receiving SGLT2 inhibitors. Before initiating, assess volume status and renal function in patients with impaired renal function (eGFR less than 60 mL/min/1.73 m<sup>2</sup>), elderly patients, patients with low systolic blood pressure, or patients on loop diuretics. In patients with volume depletion, correct this condition. After initiating, monitor for signs and symptoms of volume depletion and renal function.

### **Urosepsis and Pyelonephritis**

Serious urinary tract infections including urosepsis and pyelonephritis requiring hospitalization have been identified in patients receiving SGLT2 inhibitors, including BRENZAVVY. Treatment with BRENZAVVY increases the risk for urinary tract infections. Evaluate patients for signs and symptoms of urinary tract infections and treat them promptly.

### **Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues**

Insulin and insulin secretagogues (e.g., sulfonylureas) are known to cause hypoglycemia. BRENZAVVY may increase the risk of hypoglycemia when used in combination with insulin and/or an insulin secretagogue. A lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with BRENZAVVY.

### **Necrotizing Fasciitis of the Perineum (Fournier's Gangrene)**

Serious, life-threatening cases requiring urgent surgical intervention have been identified in postmarketing surveillance in both males and females with diabetes mellitus receiving SGLT2 inhibitors. Serious outcomes have included hospitalization, multiple surgeries, and death. Assess patients presenting with pain or tenderness, erythema, or swelling in the genital or perineal areas, along with fever or malaise. If suspected, start treatment, and discontinue BRENZAVVY.

### **Genital Mycotic Infections**

BRENZAVVY increases the risk of genital mycotic infections. Patients who have a history of genital mycotic infections or who are uncircumcised are more likely to develop genital mycotic infections. Monitor and treat appropriately.

**MOST COMMON ADVERSE REACTIONS (>5%):** Female genital mycotic infections, urinary tract infection and increased urination.

### **USE IN SPECIFIC POPULATIONS**

- Pregnancy: BRENZAVVY is not recommended during the second and third trimesters.
- Lactation: BRENZAVVY is not recommended when breastfeeding.
- Geriatric patients: There is a higher incidence of adverse reactions related to volume depletion.
- Renal Impairment: There is a higher incidence of adverse reactions related to reduced renal function.
- Hepatic Impairment: BRENZAVVY is not recommended for patients with severe hepatic impairment.

### **DRUG INTERACTIONS:**

Inducers of UGT1A9 could result in more rapid clearance of BRENZAVVY by metabolism. Doses of insulin and sulfonylureas may need to be reduced to offset the action of BRENZAVVY. The safety of BRENZAVVY is compromised when it is coupled with insulin or an insulin secretagogue (sulfonylureas and meglitinides – the latter rarely used in the US). Lithium carbonate is used as a mood stabilizer in bipolar disorder. Lithium ions might be preferentially (compared to sodium ions) taken up with glucose in the kidney. Empirical evidence has shown that lithium levels can be lower when SGLT2 inhibitors are administered. SGLT2 inhibitors produce pronounced glucosuria, which makes urine testing for glucose diagnostically useless. Measurements of 1,5 anhydroglucitol are also compromised.

*For additional important safety information about BRENZAVVY, please see the full [Prescribing Information](#).*

## **About TheracosBio**

TheracosBio develops novel therapeutics for diseases with significant societal impact. The mission of TheracosBio is to expand access to new medications for patients with common diseases.

## **About Mark Cuban Cost Plus Drug Company**

The Mark Cuban Cost Plus Drug Company, PBC aims to fundamentally change the way the pharmaceutical industry operates. As a public-benefit corporation, its social mission of improving public health is just as important as the bottom line. Cost Plus Drugs transparently charges a standard markup on every drug it sells. The [costplusdrugs.com](http://costplusdrugs.com) online pharmacy launched in January 2022 now carries over 2,000 generic and brand prescription products, delivered by mail to thousands of happy customers every day. In 2024, Cost Plus Drugs launched its Cost Plus Drugs Marketplace to extend drug savings to healthcare business purchasers.

## **Media Contact**

Brian Reid for TheracosBio  
[brian@reidstrategic.com](mailto:brian@reidstrategic.com)